



University of Southern Denmark

**Internet-based treatment of anxiety and depression in patients with ischaemic heart disease attending cardiac rehabilitation  
a feasibility study (eMindYourHeart)**

Helmark, Lotte; Ahm, Robert; Andersen, Christina Maar; Skovbakke, Søren Jensen; Kok, Robin Niels; Wiil, Uffe Kock; Schmidt, Thomas; Hjelmberg, Jacob v. B.; Frostholm, Lisbeth; Frydendal, Ditte H; Hansen, Tina Birgitte; Zwisler, Ann Dorthe Olsen; Pedersen, Susanne S.

*Published in:*

European Heart Journal - Digital Health

*DOI:*

10.1093/ehjdh/ztab037

*Publication date:*

2021

*Document version:*

Final published version

*Document license:*

CC BY-NC

*Citation for pulished version (APA):*

Helmark, L., Ahm, R., Andersen, C. M., Skovbakke, S. J., Kok, R. N., Wiil, U. K., Schmidt, T., Hjelmberg, J. V. B., Frostholm, L., Frydendal, D. H., Hansen, T. B., Zwisler, A. D. O., & Pedersen, S. S. (2021). Internet-based treatment of anxiety and depression in patients with ischaemic heart disease attending cardiac rehabilitation: a feasibility study (eMindYourHeart). *European Heart Journal - Digital Health*, 2(2), 323–335. <https://doi.org/10.1093/ehjdh/ztab037>

Go to publication entry in University of Southern Denmark's Research Portal

**Terms of use**

This work is brought to you by the University of Southern Denmark.

Unless otherwise specified it has been shared according to the terms for self-archiving.

If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim. Please direct all enquiries to [puresupport@bib.sdu.dk](mailto:puresupport@bib.sdu.dk)

# Internet-based treatment of anxiety and depression in patients with ischaemic heart disease attending cardiac rehabilitation: a feasibility study (eMindYourHeart)

Charlotte Helmark <sup>1,2\*</sup>, Robert Ahm<sup>2</sup>, Christina M. Andersen<sup>2</sup>, Søren J. Skovbakke<sup>2</sup>, Robin Kok<sup>2</sup>, Uffe K. Wiil<sup>3</sup>, Thomas Schmidt<sup>3</sup>, Jacob Hjelmberg<sup>4</sup>, Lisbeth Frostholt<sup>5</sup>, Ditte H. Frydendal<sup>5</sup>, Tina B. Hansen<sup>1</sup>, Ann-Dorthe Zwisler<sup>6,7</sup>, and Susanne S. Pedersen<sup>2,6,8</sup>

<sup>1</sup>Department of Cardiology, Zealand University Hospital, Vestermarksvej 21, DK-4000 Roskilde, Denmark; <sup>2</sup>Department of Psychology, University of Southern Denmark, Campusvej 55, DK-5230 Odense, Denmark; <sup>3</sup>The Maersk Mc-Kinney Moller Institute, University of Southern Denmark, Campusvej 55, DK-5230 Odense, Denmark; <sup>4</sup>Epidemiology, Biostatistics and Biodemography, University of Southern Denmark, J.B. Winslows Vej 9, DK-5000 Odense C, Denmark; <sup>5</sup>Research Clinic for Functional Disorders and Psychosomatics, Aarhus University Hospital, Palle Juul-Jensens Boulevard 99, DK-8200 Aarhus N, Denmark; <sup>6</sup>Department of Cardiology, Odense University Hospital, J.B. Winslows Vej 4, DK-5000 Odense, Denmark; <sup>7</sup>Danish Knowledge Center for Rehabilitation and Palliative Care (REHPA), Odense University Hospital and University of Southern Denmark, Vestergade 17, DK-5800 Nyborg, Denmark; and <sup>8</sup>Open Patient data Explorative Network (OPEN), Odense University Hospital, J.B. Winslows Vej 9, DK-5000 Odense C, Denmark

Received 12 February 2021; revised 25 March 2021; editorial decision 16 March 2021; accepted 6 April 2021; online publish-ahead-of-print 6 April 2021

## Aims

Anxiety and depression are prevalent in 20% of patients with ischaemic heart disease (IHD); however, treatment of psychological conditions is not commonly integrated in cardiac rehabilitation (CR). Internet-based psychological treatment holds the potential to bridge this gap. To examine the feasibility of an eHealth intervention targeting anxiety and depression in patients with IHD attending CR.

## Methods and results

We used a mixed-methods design, including quantitative methods to examine drop-out and change in anxiety and depression scores, and qualitative methods (thematic analysis) to evaluate patients' and nurses' experiences with the intervention. The therapist-guided intervention consisted of 12 modules provided via a web-based platform. The primary outcome was drop-out, with a drop-out rate <25% considered acceptable. Patients were considered as non-drop-out if they completed  $\geq 5$  modules. Out of 60 patients screened positive for anxiety and/or depression, 29 patients were included. The drop-out rate was 24% (7/29). Patients had a mean improvement in anxiety and depression scores of 5.5 and 4.6, respectively. On average, patients had 8.0 phone calls with their therapist and 19.7 written messages. The qualitative analysis of patients' experiences identified four themes: treatment platform, intervention, communication with therapist, and personal experience. Patients were positive towards the intervention, although some found the assignments burdensome. From the nurses, we identified three themes: intervention, inclusion procedure, and collaboration with study team. The nurses were positive, however, due to limited time some struggled with the inclusion procedure.

## Conclusion

Integrating an eHealth intervention in CR is feasible and the drop-out rate acceptable.


\* Corresponding author. Tel: +DK 4732 6111, Email: [kche@regionsjaelland.dk](mailto:kche@regionsjaelland.dk)

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

## Graphical Abstract

### Internet-based treatment of anxiety and depression in patients with ischemic heart disease attending cardiac rehabilitation: The eMindYourHeart feasibility study



## Keywords

Anxiety • Cardiac rehabilitation • Depression • E-health • Internet-based cognitive behavioural therapy • Ischaemic heart disease

## Introduction

More than 83.5 million people in Europe live with cardiovascular disease, including ischaemic heart disease (IHD).<sup>1</sup> Approximately 20% of patients with IHD have symptoms of anxiety or depression that warrant treatment.<sup>2</sup> Psychological conditions such as anxiety and/or depression act as barriers to treatment adherence of cardiac rehabilitation (CR). This increases the risk of adverse outcomes in this vulnerable subset of patients,<sup>2</sup> as evidence-based CR is shown to reduce cardiovascular mortality, rehospitalization,<sup>3</sup> and improve psychological distress.<sup>4</sup>

Clinical guidelines recommend screening for psychological distress in patients with IHD combined with psychological treatment if patients screen positive.<sup>5</sup> However, psychological treatment is not routinely offered as part of standard CR following screening.<sup>6</sup> Cognitive behavioural therapy (CBT) is an effective treatment for anxiety and depression in cardiac patients,<sup>7</sup> but lack of psychologists and challenged healthcare budgets hinder implementation of CBT in CR.<sup>6,8,9</sup>

Current technology has enabled the delivery of internet-based psychological treatment and internet-based CBT (ICBT) appear to be an equally effective treatment for anxiety and depression as face-to-face therapy.<sup>10</sup> ICBT interventions are typically delivered on a weekly basis and contain structured written material, audios, and videos, resembling face-to-face treatment.<sup>11</sup> ICBT is not geographically

or timely confined, which enables patients with internet access to engage in the treatment from any place at any time of the day.<sup>11</sup> Further advantages include reducing delay to psychological treatment, enabling quick access to treatment.<sup>11</sup>

The widespread use of internet-enabled devices ensures the possibility of delivering ICBT to a broad population, including all socio-economic backgrounds.<sup>12</sup> However, implementing the service in routine care is challenging due to complex and fragmented health-care systems, lack of funding, and limited stakeholder knowledge.<sup>12</sup> Therefore, it can be challenging for patients with IHD to obtain treatment, since there—at least in Denmark—are no automatic referral channels from the CR centres for ICBT or other psychological therapies. This means that patients with IHD besides dealing with their somatic disease must contact their general practitioner for guidance and referral to psychological treatment or seek help elsewhere.<sup>2</sup> For some patients, particularly those with depression, this may be an insurmountable task, with the risk that they never receive psychological treatment. In addition, psychological treatment might only be reimbursed under specific conditions and unaffordable for patients with e.g. low socio-economic status. Hence, integration of ICBT as part of CR has a considerable potential for the subset of patients with IHD suffering from anxiety and depression.

A systematic review on ICBT in populations with a chronic disease showed significant improvements in anxiety and depressive symptoms, but further research in cardiac populations is suggested due to

limited evidence.<sup>13</sup> Furthermore, this study points at exploring how to best integrate ICBT in routine care, as most studies have been conducted in research settings.<sup>13</sup> Results on effect of ICBT targeting anxiety and depression in cardiac patients are mixed and very limited. Two studies found no effects on symptoms of anxiety and depression in randomized controlled trials (RCTs) of patients with myocardial infarction<sup>14</sup> and heart failure,<sup>15</sup> respectively, while two other RCTs show positive effects on anxiety and depression in cardiac populations,<sup>8,16</sup> indicating a potential for ICBT in cardiac care. Thus, the high prevalence of anxiety and depression highlights the need to develop and evaluate the potential efficacy of ICBT models in CR.

The overall objective of the study was to examine the feasibility of an eHealth intervention targeting anxiety and depression in patients with IHD and integrated in standard CR as a precursor to an RCT.<sup>17</sup> Focus areas of the study were acceptability from patients and CR nurses to assess how the intervention were received and implementation of the intervention to test logistics before the RCT.<sup>18</sup> The specific aims were to evaluate (i) the dropout rate to assess acceptability, (ii) changes in anxiety and depression scores pre- and post-intervention to determine potential benefits and harms of the intervention, (iii) the extent of use of the treatment platform, (iv) the utility and experiences of the intervention from the patient's perspective, and (v) the logistics of the intervention from the CR nurses perspective.

Apart from dropout rate, the evaluations did not include predefined outcome targets as these parameters were conducted as proof of concept before the RCT.

## Methods

### Design

This feasibility study is part of the eMindYourHeart study<sup>17</sup> and was conducted using a prospective mixed-methods design to enhance the feasibility assessment of the intervention and the procedures.<sup>19</sup> The use of parallel research methods was chosen to obtain a broader view of the context of the study and to draw on strengths and mitigate weaknesses from each individual method. Quantitative methods included descriptive analysis of dropout, changes in anxiety and depression scores post-intervention (i.e. at 3 months of follow-up) and the extent of use of the treatment platform. Qualitative methods included thematic text analysis of written evaluations from patients about their experiences with the intervention and from CR nurses' their experiences with the inclusion procedures. The study is reported according to the CONSORT 2010 guideline (Consolidated Standards of Reporting Trials), using the checklist for feasibility trials.<sup>20</sup>

### Participants

#### Participants in the intervention

We aimed to recruit 30 participants, which was a pragmatic choice and considered sufficient for the aims of the feasibility evaluation. Participants were Danish patients attending CR who screened positive on the Hospital Anxiety and Depression Scale (HADS)<sup>21</sup> as part of routine care. Inclusion criteria were age  $\geq 18$  years, diagnosis of IHD, HADS score  $\geq 8$  on depression and/or anxiety,<sup>22</sup> access to a computer or smartphone, ability to use computer or smartphone, proficient in the Danish language, and a signed informed consent. Exclusion criteria were severe psychiatric disorder (e.g. schizophrenia), severe cognitive difficulties (e.g. dementia), participation in other psychological intervention studies, seeing a

psychologist or mental health professional for the treatment of depression and anxiety.

The participants were recruited consecutively from nine Danish CR centres. The centres were strategically chosen across all five Regions in Denmark and across hospitals and municipalities, to ensure variation with respect to logistics in the CR programmes and a variety in patient demographics. CR nurses recruited patients for the study when starting their CR programme and being routinely screened for anxiety and depression, as recommended by national<sup>23</sup> and European guidelines.<sup>5</sup>

#### Participants in the evaluations

All patients who completed the intervention and all CR nurses involved in recruitment of patients were invited to participate in an evaluation of the intervention.


### Intervention

The intervention was developed using a participatory design.<sup>24</sup> This design was used to ensure the target patient group experienced the ICBT as relevant and useful. Psychologists and supervised graduate students of psychology (in their final year) with thorough training in mental health in cardiac patients guided the intervention as therapists. A CR nurse supervised the cardiac aspects when needed. The therapist contacted the patients within three workdays after completing the baseline questionnaire. The intervention began with a diagnostic telephone interview conducted by the therapist, using a brief purpose-designed interview protocol. The protocol was implemented to gain background information about patients' life situation, experiences with heart disease, motivation to participate in the intervention, understanding of the concepts of the ICBT intervention, sleep problems, use of alcohol and recreational drugs, previous experiences with anxiety, depression and psychological treatment, suicidal ideation, and prior trauma as well as building a therapeutic alliance. The 12-week ICBT intervention consisted of nine mandatory and three voluntary modules, covering topics like behavioural activation and cognitive restructuring (Figure 1; Table 1). All mandatory modules contained written assignments. Patients could access the treatment platform at any time of the day using smartphone, tablet, or computer.

Patients had a personal therapist assigned who supported them throughout the intervention, guiding them through the modules on a weekly basis by asynchronous written messages via the platform. Patients could use the written message function whenever they wanted, and the therapist would respond within two workdays. Furthermore, patients had four phone calls with the therapist during the intervention, including the diagnostic interview. In case patients had an additional need, further phone calls could be assigned.<sup>25</sup> The dropout rate was tried diminished by therapists supporting all patients with individual approaches depending on the patient's situation and the phone calls allowed for this individualization. Patients accessed the intervention platform by a GDPR-compliant website, using their NemID, a Danish national two-factor authentication solution.

### Quantitative outcomes and statistics

The primary outcome was dropout rate where a rate  $<25\%$  was considered a success, based on a systematic review of guided ICBT.<sup>26</sup> Treatment completion was defined a priori as five or more mandatory modules completed, since this would ensure that the patient had engaged in both psycho-educative and cognitive and behaviour change content. Secondary outcomes were changes in HADS scores, extent of use of the platform and utility of intervention. Utility was evaluated with six generic questions from the Internet Evaluation and Utility Questionnaire,<sup>27</sup> which were chosen as they fitted well with the context of the current intervention. Data on dropout and patient-reported outcomes were collected



and managed electronically using the internationally recognized system for data management—Research Electronic Data Capture (REDCap)—via Odense Patient data Explorative Network (OPEN).<sup>28</sup> Patient-reported data were collected at baseline and at 3-month follow-up. Extent of use of the platform included duration spent on the treatment platform and frequency and time of login. These data were captured via the treatment platform. Demographics were collected in the baseline questionnaire and clinical data through the Danish Cardiac Rehabilitation Database.<sup>29</sup> Results were presented using descriptive statistics and reported with frequencies, percentages, means, range, and standard deviation where appropriate. All statistical analyses were conducted using Stata version 16.1.

### Qualitative measurements

Data on patients' experiences with the intervention were generated through a purpose-designed questionnaire at 3 months of follow-up, divided into categories regarding experiences with the intervention. Patients gave written feedback on the questionnaire, which did not call for alterations. Data on CR nurse's perception of the intervention were likewise generated through a purpose-designed questionnaire divided into categories regarding logistics of the recruitment procedure. Two CR nurses gave written feedback on this, leading to minor adjustments. Both questionnaires were completed online using REDCap, and answered with open text, enabling the possibility of broad feedback. Both groups had the possibility to write issues beyond the predefined categories and give their evaluation by phone if preferred. The qualitative analysis processes were performed using thematic analysis, inspired by Braun and Clarke.<sup>30</sup> For each evaluation, two authors (C.H. and S.J.S.) repeatedly read and coded the data independently and afterwards discussed the coding until consensus was reached. The codes were then sorted into categories and mapped into themes. To optimize credibility the themes were discussed, analysed, and revised with the inclusion of further two authors (C.M.A. and S.S.P.). The trustworthiness of the qualitative analysis was tried accomplished by involving the end-users in the design of the questionnaires to optimize reliability, by using investigator triangulation in the coding process to gain credibility and by describing the context to show transferability.

### Hospital Anxiety and Depression Scale (HADS)

The HADS was chosen as the screening tool for anxiety and depression, since it has shown to be a valid measure in cardiac patients<sup>22,31</sup> and is recommended in CR in the Danish national clinical guidelines.<sup>23</sup> HADS is a 14-item questionnaire with 7 items each contributing to the subscales anxiety and depression. Each item is scored on a Likert scale from 0 to 3 leading to a score range from 0 to 21 for both subscales, with higher scores indicating more severe symptoms. The commonly used cut-off score of  $\geq 8$  was applied to identify symptoms of anxiety and depression.<sup>22</sup>

### Ethics

Permission to conduct the study was obtained from the Danish Data Protection Agency at Odense University Hospital (17/41433 on 24 November 2017) via the umbrella permission of the Region of Southern Denmark. Ethical approval was obtained from the Regional Committees on Health Research Ethics for Southern Denmark (S-20180024). The study complies with the Helsinki Declaration with all patients providing written informed consent.<sup>32</sup>

## Results

### Participants and drop-out rate

The participants were recruited and treated between November 2019 and July 2020. *Figure 2* outlines the flowchart of the study population. A total of 60 patients were identified with elevated HADS scores, of which six were excluded based on exclusion criteria, while 16 patients declined to participation and one died. After the baseline questionnaire and telephone interview, a further eight were excluded or declined. Thus, we included 29 participants in the feasibility study corresponding to 54% of eligible patients. Of the included patients, five dropped out and two did not complete the required five modules to be considered 'completer' of the intervention. This means we had a dropout rate of 24.1% (7/29), reaching the desired rate of <25%.

**Table 1** Description of the modules in the internet-based cognitive behavioural treatment

Module	Content	Aim	Therapeutic components
Extra	Overview of treatment	Introduction to treatment + information about IHD.	Psychoeducation
1	Information and expectations about treatment	To give the patients a better understanding of their own psychological reactions.	Psychoeducation, motivation, and goals
2	How to handle difficult emotions	To help patients understand and manage their own emotions in a healthier way.	Psychoeducation, mentalization, self-compassion
3	Cognitive change (Introduction to cognitive diamond)	To give patients a better understanding about their own psychological patterns and concrete tools on how to manage difficult thoughts.	Psychoeducation, cognitive restructuring, cognitive diffusion
4	Attention training and control and avoidance and behaviour	To help patients to less attention on bodily anxiety symptoms which can be interpreted as IHD. Moreover, to help them identify their own safety and avoidance behaviours.	Psychoeducation, selective attention training
5	Exposure therapy	To help patients face their fears and overcome them.	Exposure therapy
6	Internal barriers, balancing own efforts in treatment and self-care	Patients identify and work with potential treatment barriers. Also benefits of self-compassion and exercises promoting self-compassion.	Cognitive restructuring, self-compassion
7	Behavioural activation and exercise in daily life	To activate depressive patients and to promote physical exercises, to lower depression and anxiety.	Behavioural activation
8	Acceptance of heart condition and finding life values	To promote acceptance of current situation, and a shift in focus away from limitations and towards possibilities.	Cognitive restructuring, life values
9	Relapse prevention	To ensure patients know how to spot early signs of relapse and know how to handle these.	Psychoeducation, cognitive restructuring
Extra	Sleep	Knowledge and tools on how to improve sleep.	Psychoeducation
Extra	Lifestyle changes	Information about living with IHD, e.g. where to get help quitting smoking.	Psychoeducation

Out of 54 eligible patients, 25 (46%) were not included in the study. For these 25 patients, data are missing for five since they did not provide informed consent. Of the remaining 20 patients, four patients (20%) were female, and the mean age was 59.7 (SD: 10.1), with a range of 33–76 years. The characteristics of the 29 included patients are presented in [Tables 2](#) and [3](#).


### Changes in anxiety and depression (measured with HADS)

We found a mean improvement for HADS anxiety scores at 5.5 points and a mean improvement for HADS depression scores at 4.6 points ([Table 4](#)). Among the 22 completed patients, three had missing HADS scores at follow-up. The minimal clinical important difference (MCID) reflects meaningful changes for the patient, and for HADS scores it is 1.8.<sup>33</sup> We found that 16 out of 19 (84%) patients achieved the MCID for anxiety and 15 out of 19 patients (79%) achieved the MCID for depression. Three reported worse outcomes post-

intervention compared to pre-intervention, where one of these were clinically relevant according to the MCID.

### Use of intervention

Patients who completed the intervention had a mean of 8 telephone calls and 19.7 written messages with their therapist during the intervention ([Table 5](#)). The diagnostic telephone interview is not included in these data. On average, patients spent 10.95 h on the treatment platform. Participants were automatically logged out after 5 min of inactivity, so the time consumption does not include patients who left the treatment platform unattended. For patterns of login, we found that patients logged on to the treatment platform a mean of 33.5 times, where 59.4% of the times took place during normal opening hours for CR centres, while 40.6% took place outside normal opening hours. The total time of the intervention from diagnostic interview to completion of the last module was on average 13.3 weeks.



## Utility of the intervention

The same 19 patients who completed HADS also completed the utility questions post-intervention (Figure 3). The majority found the intervention easy to use, that the intervention kept their interest and attention, that the information was useful and easy to understand and were satisfied with the intervention. The majority were not worried about their privacy although one patient-reported being worried 'mostly'.

## Patients' evaluation of the intervention

Of the 22 patients who completed the intervention, 14 agreed to provide an evaluation. The text analysis led to the identification of four themes: treatment platform; intervention; communication with therapist; and personal experience (Table 6). Patients experienced that accessing and navigating on the treatment platform was easy, however few found the logon procedure burdensome. Regarding the intervention, the patients were highly satisfied with the concept,

**Table 2** Characteristics of included patients (n = 29)

	Completed participants (n = 22)	Non-completed participants (n = 7)
Age, mean (SD)	64.6 (11.2)	53.0 (10.0)
range	47–84	40–70
Female sex, n (%)	9 (41%)	2 (29%)
Living alone, n (%)	6 (27%)	4 (57%)
Employment status, n (%)		
Employed	9 (41%)	5 (71%)
Retired	12 (55%)	1 (14%)
Other	1 (4%)	1 (14%)
Highest completed education, n (%)		
Primary school/high school	3 (14%)	2 (29%)
Short education (≤3 years)	7 (31%)	5 (71%)
Bachelor's degree (3–4 years)	9 (41%)	0
Higher education (≥5 years)	3 (14%)	0
Body mass index, mean (SD)	28.1 (5.3)	30.2 (4.6)
range	20.6–40.9	23.3–37.2
Currently smoking, n (%)	2 (9%)	3 (43%)
Previously diagnosed with psychiatric disorder, n (%)	2 (9%)	1 (14%)
Physically active ≥150 min per week, n (%)	7 (32%)	2 (29%)

**Table 3** Clinical data from the Danish Cardiac Rehabilitation Database (n = 18)<sup>a</sup>

Variable	n (%)	Missing n (%)
Primary indication for referral to cardiac rehabilitation		
ST-elevation myocardial infarction	5 (28)	2 (11)
Non-ST-elevation myocardial infarction	7 (39)	
Stable angina pectoris	4 (22)	
Primary cardiac procedure		
Percutaneous coronary intervention	11 (61)	0 (0)
Medically managed <sup>b</sup>	6 (33)	
Heart valve replacement	1 (6)	
Diabetes	1 (6)	0 (0)
Heart failure	1 (6)	0 (0)
Referred for exercise-based cardiac rehabilitation	16 (89)	1 (6)
Referred for group-based patient education	11 (6)	2 (11)

SD, standard deviation.

<sup>a</sup>Out of the 22 completed participants, 18 were identified in the Danish Cardiac Rehabilitation Database.<sup>b</sup>These are classified as medically managed, as they are not registered as having undergone percutaneous coronary intervention or coronary artery bypass grafting.

leaving them in control of time, place, and speed. Still, some found the written assignments inside some of the modules burdensome. All patients were highly satisfied with the communication with the therapist, appraising the competences of their personal therapist. The patients indicated that their personal motivation for participation was related to a need for help with worries, fear, anxiety, and loneliness, while some specified tools for coping. They experienced the

**Table 4** Changes in anxiety and depression (measured with HADS) (n = 19)<sup>a</sup>

	HADS pre-scores	HADS post-scores	Change
HADS A			
Mean (SD)	11.7 (3.1)	6.1 (4.4)	<b>−5.5 (4.6)</b>
Range	6–18	0–15	-16 to +4
HADS D			
Mean (SD)	9.5 (3.2)	4.9 (3.8)	<b>−4.6 (5.1)</b>
Range <sup>b</sup>	5–17	0–12	-15 to +5

HADS, Hospital Anxiety and Depression Scale; SD, standard deviation.

<sup>a</sup>Missing HADS data at follow-up, n = 3.<sup>b</sup>Reported worse outcomes post-intervention, n = 3.

intervention helped with better understanding and acceptance of difficult feelings, and the majority pointed at the cognitive diamond as particularly helpful. Some were positively surprised at the effectiveness of online intervention, and some found it a relief not having to engage in face-to-face treatment. The patients had individual patterns for when they engaged in the intervention based on energy levels, moods, convenience, and schedules.

### Nurses' evaluation of the intervention

Among the involved CR nurses, 14 completed the evaluation, representing all nine centres. The text analysis led to three themes: intervention; inclusion procedures; and collaboration with study team (Table 7). Overall, the nurses were very positive towards the concept of ICBT, appreciating fast access to psychological treatment for their patients. The nurses experienced that patients do not want to or have the energy to contact their general practitioner seeking help for



**Table 5 Use of intervention for completed participants (n = 22)**

	Mean (SD)	Range
Number of contacts with therapist		
Telephone calls	8.0 (3.9)	3–22
Written messages on the treatment platform <sup>a</sup>	19.7 (22.6)	5–37
Time consumption on the treatment platform (h)	10.95 (6.5)	4.7–34.1
Number of log ons to the treatment platform <sup>b</sup>		
Total (100%)	33.5 (13.4)	12–69
Logins within normal opening hours <sup>c</sup> (59.4%)	19.9 (8.3)	4–34
Logins outside normal opening hour <sup>c</sup> (40.6%)	13.6 (7.3)	3–35
Total time of intervention including diagnostic interview (weeks)	13.3 (2.8)	10–20

SD, standard deviation.

<sup>a</sup>Combined written messages on the treatment platform from participant and therapist.

<sup>b</sup>Including training of participant, which included logging on twice.

<sup>c</sup>Opening hours defined as Monday to Friday from 8 a.m. until 4 p.m., public holidays excluded.

their anxiety and depression. They also experienced that some patients were not interested in online treatment, and for others it was offered to soon in the CR pathway. Regarding inclusion procedures, some were challenged by the additional task of including patients as part of clinical practice, but also that procedures gradually became less burdensome. Willingness to use clinical time on recruiting patients was in recognition of lack of psychological treatment within their CR centre. Sustainable support from the study team was highly appreciated, safeguarding motivation to participation in the study.

## Discussion


This study examined the feasibility of integrating ICBT into routine CR in patients with IHD. We found a dropout rate from the ICBT

intervention of 24%, which is below the a priori determined threshold. The study showed satisfactory improvements in anxiety and depression scores during the intervention. User patterns of the intervention showed a large variability. The evaluations revealed that patients were very positive towards the intervention, but some experienced the workload burdensome. The nurses were very positive about the concept of ICBT but stated that including patients was yet another task to manage in busy clinical practice.

## Acceptability and adherence

Among 54 eligible patients with high HADS scores, 29 were included in the study. Of the 25 eligible non-included patients, six experienced minor mental problems, four were excluded at baseline with HADS scores  $\leq 8$ , 1 died, leaving 14 out of 54 eligible patients (26%) with symptoms of anxiety and depression who declined participation. Three of these patients reported lack of energy and for four patients we do not have information about reason for declining. We can only speculate that for some patients the timing of the intervention may have been too soon after the cardiac event. Others may be unwilling to engage in a psychological intervention e.g. due to stigma or being uncomfortable with online treatment. To our knowledge, only one other study on ICBT within cardiology included patients in routine care, and they reported 1359 out of 1946 eligible patients (70%) declined participation.<sup>14</sup> This indicates that this study has a satisfactory acceptability. To safeguard inclusion, additional nine CR centres are recruited for the RCT.

Only few studies on ICBT within cardiology are published. When comparing our findings with these, one RCT in patients with myocardial infarction found no effect of ICBT on symptoms of anxiety or depression.<sup>14</sup> The study had a low treatment adherence with 38.4% completing only the introductory module and 15.4% completing additional modules, making it difficult to evaluate the effect of the intervention. Another RCT of ICBT in patients with cardiovascular disease found a significant effect on depression and had 59 patients (82%) completing at least four out of seven modules.<sup>8</sup> A recent RCT in patients with acute coronary event also showed a positive effect of ICBT on both anxiety and depression where 92% completed at least four out of eight lessons.<sup>16</sup> This suggests that poor adherence to



**Figure 3** Utility of the intervention.

**Table 6** Summary of patient's experiences based on the written evaluations

Theme	Findings	Quotes
Treatment platform	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Easy to access, navigate, and use</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• Burdensome logon procedure</li> <li>• Miss possibility to follow progress in the modules and to print out pages</li> </ul>	<p>'Nice, easy and clear. Miss possibility to print out on paper' (female, 73 years)</p>
Intervention	<p>Pro:</p> <ul style="list-style-type: none"> <li>• In control of time and speed of the intervention</li> <li>• Very relevant content</li> <li>• Highly satisfied with the concept</li> <li>• Can participate from home</li> <li>• Fine balance between treatment elements</li> <li>• Length of intervention and amount of homework appropriate</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• Length of time to intervention too short</li> <li>• Amount of homework too much</li> </ul>	<p>'With online (treatment)—I decide for myself when to engage in the treatment' (female, 73 years)</p> <p>'It was nice that I did not have to leave home even though I needed help to live without anxiety' (male, 60 years)</p> <p>'Written assignments were a bit burdensome, especially when you have limited energy' (female, 52 years)</p>
Communication with therapist	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Highly satisfied with therapist</li> <li>• Therapist is competent</li> <li>• Sufficient time in phone calls</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• None</li> </ul>	<p>'100% okay. It is a really good idea that you can text with your therapist during the intervention' (female, 79 years)</p>
Personal experience	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Positively surprised of the concept</li> <li>• More relaxed and calm afterwards</li> <li>• Better acceptance of feelings</li> <li>• Better understanding of own reactions</li> <li>• Obtained various tools to cope with negative thoughts and feelings</li> <li>• No physical contact with therapist</li> <li>• Cognitive diamond helpful</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• Not enough time to go through the modules</li> <li>• Too many written assignments</li> </ul>	<p>'I liked to see my psychologist in person, which I have tried before. But I am positively surprised that this (online treatment) can work' (female, 64 years)</p> <p>'Actually this (online treatment) is an advantage when you are a bit of an "introvert" like me. I did not have to look anybody in the eyes' (male, 56 years)</p>

treatment is a barrier to dissemination of potentially effective treatments, even though a linear relationship is still fully uncovered<sup>34</sup> and the association was insignificant in the RCT by Lundgren et al.<sup>15</sup> It has

been proposed from previous ICBT studies that a factor attributing to poor adherence might be a lack of customization of the ICBT treatment to fit the target population.<sup>8,16,35</sup> This is among the reasons

**Table 7** Summary of nurses' experiences based on the written evaluations

Theme	Findings	Quotes
Intervention	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Online treatment is a valuable and important concept</li> <li>• Most patients were interested and positive</li> <li>• Free of charge for the patients</li> <li>• Fast track to psychological treatment</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• Some patients had doubts or rejected the offer because treatment is online</li> <li>• Some patients might need the treatment later in the care pathway</li> </ul>	<i>'The best part is to have an offer to our sad patients, and on top of that, that the patients can start the intervention so fast'</i>
Inclusion procedures	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Good introduction</li> <li>• Inclusion process became easier with routine</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• Challenging on top of usual assessment tasks at start of CR</li> <li>• Too much project information to both patients and the nurses</li> </ul>	<i>'Good project, but we also have to register in other databases, leaving less time for the patient'</i>
Collaboration with study team	<p>Pro:</p> <ul style="list-style-type: none"> <li>• Personal introduction to the study</li> <li>• High level of information, feedback and availability of the team</li> </ul> <p>Con:</p> <ul style="list-style-type: none"> <li>• None</li> </ul>	<i>'The best part is to feel that someone is passionate about this (psychological treatment)'</i>

why we decided using a participatory design for development of the intervention.<sup>24</sup> There are substantial differences in recruitment procedures, length of interventions as well as numbers and content of the modules in the mentioned studies, making comparisons difficult. In the current study, patients were considered 'completers' if they had completed at least five of the mandatory modules. Two patients did not accomplish the required five modules due to personal life situations but still benefitted from the intervention based on the treatment evaluation of their therapist, suggesting that our choice of five modules might be somewhat arbitrary. On these grounds, and since there is no gold standard for what comprises a 'completer' in ICBT, we have decided to change our definition in the eMindYourHeart RCT. Patients will be considered completers regardless of number of completed modules, if they follow the treatment plan agreed upon with their therapist. A patient can drop out of the treatment in two ways: (i) active drop out by communicating this decision to their

therapist, or (ii) passive drop out, i.e. the patient stops responding to their therapist attempts to contact them and the contact is not re-established within the 12 weeks of treatment. We have extended the treatment manual and developed a stricter, therapist independent rating system for assessing if a module is completed, and which modules are relevant for the individual patient. This to secure a more personalized approach depending on the patient's individual situation and a high degree of interrater reliability.

### Changes in anxiety and depression

Given the design of this uncontrolled study, we did not analyse the effectiveness of the intervention. With mean improvements in scores of 5.5 for anxiety and 4.9 for depression, the intervention indicated positive signs of obtaining the established MCID of 1.8.<sup>33</sup> Although these results could be due to chance, they are encouraging. At an individual level, 18/22 (82%) of those patients who completed the

intervention reached the cut-offs for absence of anxiety and depression (HADS scores < 8). Schneider *et al.*<sup>16</sup> showed a range of recovery from anxiety and depression at 77%-85% that resembles our results. Regarding the three patients who had worsened HADS scores post-intervention, two had their treatment prolonged. Therefore, their follow-up HADS questionnaire was filled out before the intervention was completed, and in both cases, their therapist assessed clinical improvement at the end of the intervention. The third patient was severely impacted by COVID-19 and the associated social isolation. Out of the 22 completed patients, three stopped filling out the follow-up battery of questionnaires before they reached the HADS questionnaire, despite receiving two reminders. Since HADS will be the primary outcome of the eMindYourHeart RCT, HADS will be moved to the top of the follow-up questionnaire to safeguard this measure. In addition, the therapist will systematically encourage the patients to respond to this questionnaire at their last contact.

### Use of intervention

Regarding use of intervention, Lundgren *et al.*<sup>15</sup> found that number of logins was associated with improvements in depression in the ICBT group. However, data on numbers of logins and range are not reported making comparison difficult. Our results somewhat resemble the results of Schneider *et al.*<sup>16</sup> in their RCT, as mean numbers of login were 33.5 (SD 13.4) vs. 26.4 (SD 16.6) and mean numbers of phone calls 8.0 (SD 3.9) vs. 4.8 (SD 3.1). The number of phone calls in this study was higher than expected which could be explained by the study design, testing, and adjusting procedures. Another explanation is the start of the COVID-19 pandemic in the trial period, which induced extra calls from patients. Our analysis showed that 40.6% of logins happened outside normal opening hours of CR services, which emphasizes that patients might prefer treatment at other timepoints than currently offered. One explanation could be that negative thoughts and emotions intrude at night-time when there is more time to think and ruminate. Concerning the duration of intervention, we found an average of 13.3 weeks that is longer than the stipulated 12 weeks. One reason is delay in time for some patients between the diagnostic interview and the start of first module, mostly caused by technical issues. Moreover, some patients requested extra time to complete the intervention due to personal situation. In the RCT, the length of the intervention will be monitored from start to end of the modules and the time kept to 12 weeks, making comparison easier.

### Patient evaluation

A relatively high proportion of patients had a high educational level. We can only speculate if patients with low levels of education may be more reluctant to participate in ICBT or if CR staff are more reluctant to include patients with low levels of education, since this is seen in previous studies.<sup>16,36</sup> Since all included patients with high levels of education completed the current intervention, it could also mean that our intervention might not be properly adjusted to participants with lower levels of education, despite efforts in the participatory design study.<sup>24</sup> Some patients found the written assignments burdensome, signalling the need for personalized levels of the modules, which could potentially be related to level of education. Challenges with burden of assignments are also described previously.<sup>36,37</sup> Wallin

*et al.*<sup>36</sup> argue for a more personalized ICBT approach, as some patients perceive text-based material strenuous and time-consuming. We have accordingly adjusted the one-size-fits-all approach to a more personalized model in the following RCT. Also consistent with the findings from Wallin *et al.*,<sup>36</sup> some of our patients found the two-step logon procedure burdensome but given the GDPR rules, this is the best possible solution momentarily. Conversely, it might give other patients a feeling that the platform is secure and trustworthy, which is previously reported.<sup>37</sup> Some patients expressed a preference to print out material from the platform. Previous studies have described this,<sup>12</sup> and e.g. the Mindspot Clinic in Australia are sending out hard copies of the material upon request.<sup>38</sup> Although this will not be possible in the RCT, since it will refrain us from collecting data on user activity on the platform, it should be considered if implementing into routine CR. Most patients found the intervention easily accessible and manageable and all were highly satisfied with their personal therapist and reported positive personal gains related to their mental condition. As one wrote: 'I have become myself again, this is pure happiness'.

### Nurses evaluation

The findings from the nurses' evaluations revealed strong support for the concept of ICBT integrated into CR, as this gives patients fast access to psychological treatment. They appreciated the high level of availability of the eMindYourHeart study team, which might have a positive impact on patient recruitment. Still, it is evident that the CR nurses have a high workload and they struggled with inclusion procedures as it takes clinical time from their patients, which suggests that implementation is a complex process.<sup>39</sup> It is therefore necessary to support the nurses to keep them engaged in recruiting patients, and in case of future implementation in routine care, a solid implementation plan is needed.

The European Society of Cardiology supports digital health as an innovative opportunity to improve the quality of care in secondary prevention.<sup>9</sup> Especially in the light of the COVID-19 pandemic, ICBT might be an effective solution to treat anxiety and depression in patients with IHD as an integrated part of CR.<sup>40</sup> Patients expressed high levels of gratefulness for the continuity of this intervention and the possibility to talk about their concerns related to COVID-19, while experiencing most elements of CR being cancelled. We managed to adapt the intervention to help also with concerns related to COVID-19, as it became obvious that this induced anxiety and worries among the patients.

### Strengths and limitations

Strength of this study include the multicentre approach, testing of inclusion procedures in a large variety of CR models and centres across Denmark, and adaptations and optimisation of the intervention. Collaboration with the recruiting centres prior to the RCT also included a webinar for the CR staff (on their request) which is likely to be an advantage for successful recruitment in the RCT study. The study had following limitations: firstly, there were 16 patients declining to participate in the study, and since we only have sparse data on these patients, we are not able to examine if they differ from participating patients. Thus, there might be unidentified subgroups not willing to participate in ICBT. Declining patients are asked to fill out an

informed consent to use of data in the RCT and due to the considerably higher volume in this study, these data might reveal important knowledge. Secondly, since the feasibility study were uncontrolled the positive changes in outcomes should be interpreted with great caution. In addition, clinical data from the Danish Cardiac Rehabilitation Database were rather incomplete, and other national registries will be considered for the RCT. We asked patients and nurses to evaluate their experiences with the intervention and procedures in writing, while oral interviews potentially could have led to deeper insights into barriers and facilitators of the intervention, as done by e.g. Wallin *et al.*<sup>36</sup> However, an in-depth qualitative analysis was not the scope of this study and would have required thorough elaboration.

## Conclusion

This study demonstrated the feasibility of the eMindYourHeart study, based on drop-out rate, mean improvements in HADS scores, use of intervention and experiences from patients and CR nurses. The study was an essential prerequisite for a larger interventional study. Thus, the intervention will be adjusted according to the reported results and used in the following eMindYourHeart RCT.

## Lead author biography



Charlotte Helmark is a PhD candidate employed at Zealand University Hospital and working in the cross-field between cardiology and psychology.

## Acknowledgements

The authors would like to thank all participating patients and health-care staff involved in the study. We would also like to thank Nicolaj Knudsen and Richard Jensen from Aarhus University Hospital for their help with adjustments and data collection on the platform used for the eMindYourHeart intervention. Thank you to the Danish Cardiac Rehabilitation Database (DHRD) under the Danish Clinical Quality Program for access to clinical data.

## Funding

This trial was supported with a grant from University of Southern Denmark's Strategic funding, the Danish Heart Foundation (18-R124-A8309-22106), Region of Southern Denmark and Region of Zealand joint research foundation (A343), Odense University Hospital Research Foundation (72-A3575), Region Zealand Palliation Fund and Sygesikring Denmark (2020-0041). None of the funding agencies has had any role in neither the design of the study nor in the collection, analysis, and interpretation of the data or writing of the manuscript.

**Conflict of interest:** none declared.

## Data availability

The data from this article cannot be shared publicly due to the use of confidential health information.

## References

1. Timmis A, Townsend N, Gale C, Grobbee R, Maniadas N, Flather M, Wilkins E, Wright L, Vos R, Bax J, Blum M, Pinto F, Vardas P. European Society of Cardiology: cardiovascular disease statistics 2017. *Eur Heart J* 2018;**39**:508–579.
2. Pogossova N, Saner H, Pedersen SS, Cupples ME, McGee H, Höfer S, Doyle F, Schmid JP, von Käne R. Psychosocial aspects in cardiac rehabilitation: From theory to practice. A position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation of the European Society of Cardiology. *Eur J Prev Cardiol* 2015;**22**:1290–1306.
3. Salzwedel A, Jensen K, Rauch B, Doherty P, Metzendorf M-I, Hackbusch M, Völler H, Schmid JP, Davos C. Effectiveness of comprehensive cardiac rehabilitation in coronary artery disease patients treated according to contemporary evidence based medicine: update of the Cardiac Rehabilitation Outcome Study (CROS-II). *Eur J Prev Cardiol* 2020;**27**:1756–1774.
4. Richards SH, Anderson L, Jenkinson CE, Whalley B, Rees K, Davies P, Bennett P, Liu Z, West R, Thompson DR, Taylor RS. Psychological interventions for coronary heart disease: Cochrane systematic review and meta-analysis. *Eur J Prev Cardiol* 2018;**25**:247–259.
5. Ambrosetti M, Abreu A, Corrà U, Davos CH, Hansen D, Frederix I, Iliou MC, Pedretti RF, Schmid JP, Vigorito C, Voller H, Wilhelm M, Piepoli MF, Bjarnason-Wehrens B, Berger T, Cohen-Solal A, Cornelissen V, Dendale P, Doehner W, Gaita D, Gevaert AB, Kemps H, Kraenkel N, Laukkanen J, Mendes M, Niebauer J, Simonenko M, Zwisler AO. Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020;doi: 10.1177/2047487320913379.
6. Pedersen SS, Andersen CM. Minding the heart: why are we still not closer to treating depression and anxiety in clinical cardiology practice? *Eur J Prev Cardiol* 2018;**25**:244–246.
7. Reavell J, Hopkinson M, Clarkesmith D, Lane DA. Effectiveness of cognitive behavioral therapy for depression and anxiety in patients with cardiovascular disease: a systematic review and meta-analysis. *Psychosom Med* 2018;**80**:742–753.
8. Johansson P, Westas M, Andersson G, Alehagen U, Broström A, Jaarsma T, Mourad G, Lundgren J. An internet-based cognitive behavioral therapy program adapted to patients with cardiovascular disease and depression: randomized controlled trial. *JMIR Ment Health* 2019;**6**:e14648.
9. Frederix I, Caiani EG, Dendale P, Anker S, Bax J, Böhm A, Cowie M, Crawford J, de Groot N, Dilaveris P, Hansen T, Koehler F, Krstacic G, Lambrinou E, Lancellotti P, Meier P, Neubeck L, Parati G, Piotrowicz E, Tubaro M, van der Velde E. ESC e-Cardiology Working Group Position Paper: overcoming challenges in digital health implementation in cardiovascular medicine. *Eur J Prev Cardiol* 2019;**26**:1166–1177.
10. Andersson G, Cuijpers P, Carlbring P, Riper H, Hedman E. Guided Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: a systematic review and meta-analysis. *World Psychiatry* 2014;**13**:288–295.
11. Andersson G, Titov N, Dear BF, Rozental A, Carlbring P. Internet-delivered psychological treatments: from innovation to implementation. *World Psychiatry* 2019;**18**:20–28.
12. Titov N, Hadjistavropoulos HD, Nielssen O, Mohr DC, Andersson G, Dear BF. From research to practice: ten lessons in delivering digital mental health services. *J Clin Med* 2019;**8**:1239.
13. Mehta S, Peynenburg VA, Hadjistavropoulos HD. Internet-delivered cognitive behaviour therapy for chronic health conditions: a systematic review and meta-analysis. *J Behav Med* 2019;**42**:169–187.
14. Norlund F, Wallin E, Olsson EMG, Wallert J, Burell G, von Essen L, Held C. Internet-based cognitive behavioral therapy for symptoms of depression and anxiety among patients with a recent myocardial infarction: the U-CARE Heart Randomized Controlled Trial. *J Med Internet Res* 2018;**20**:e88.
15. Lundgren JG, Dahlström Ö, Andersson G, Jaarsma T, Köhler AK, Johansson P. Effect of guided web-based cognitive behavioral therapy on patients with depressive symptoms and heart failure: a pilot randomized controlled trial. *J Med Internet Res* 2016;**18**:e194.
16. Schneider LH, Hadjistavropoulos HD, Dear BF, Titov N. Efficacy of internet-delivered cognitive behavioural therapy following an acute coronary event: a randomized controlled trial. *Internet Interv* 2020;**21**:100324.

17. Pedersen SS, Andersen CM, Ahm R, Skovbakke SJ, Kok R, Helmark C, Wiil U, Schmidt T, Olsen KR, Hjelmberg J, Zwisler A-D, Frostholm L. Efficacy and cost-effectiveness of a therapist-assisted web-based intervention for depression and anxiety in patients with ischemic heart disease attending cardiac rehabilitation [eMindYourHeart trial]: a randomised controlled trial protocol. *BMC Cardiovasc Disord* 2021;**21**:20.
18. Bowen DJ, Kreuter M, Spring B, Cofta-Woerpel L, Linnan L, Weiner D, Bakken S, Kaplan CP, Squiers L, Fabrizio C, Fernandez M. How we design feasibility studies. *Am J Prev Med* 2009;**36**:452–457.
19. Hansen HP, Tjørnhøj-Thomsen T. Meeting the challenges of intervention research in health science: an argument for a multimethod research approach. *Patient* 2016;**9**:193–200.
20. Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, Lancaster GA; PAFS consensus group. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. *Pilot Feasibility Stud* 2016;**2**:64.
21. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;**67**:361–370.
22. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res* 2002;**52**:69–77.
23. Sundhedsstyrelsen. National Klinisk Retningslinje for Hjerterehabilitering. 2015. <https://www.sst.dk/da/udgivelser/2015/nkr-hjerterehabilitering> (10 February 2021).
24. Schmidt T, Kok R, Andersen CM, Skovbakke SJ, Ahm R, Wiil UK, Frostholm L, Pedersen SS. Development of an internet-delivered program and platform for the treatment of depression and anxiety in patients with ischemic heart disease in eMindYourHeart. *Inform Health Social Care* 2021;doi: 10.1080/17538157.2021.1878185.
25. Hadjistavropoulos HD, Schneider LH, Mehta S, Karin E, Dear BF, Titov N. Preference trial of internet-delivered cognitive behaviour therapy comparing standard weekly versus optional weekly therapist support. *J Anxiety Disord* 2019;**63**:51–60.
26. Richards D, Richardson T. Computer-based psychological treatments for depression: a systematic review and meta-analysis. *Clin Psychol Rev* 2012;**32**:329–342.
27. Ritterband LM, Ardan K, Thorndike FP, Magee JC, Saylor DK, Cox DJ, Sutphen JL, Borowitz SM. Real world use of an Internet intervention for pediatric encephalitis. *J Med Internet Res* 2008;**10**:e16.
28. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, McLeod L, Delacqua G, Delacqua F, Kirby J, Duda SN; REDCap Consortium. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019;**95**:103208.
29. Zwisler A-D, Rossau HK, Nakano A, Foghmar S, Eichhorst R, Prescott E, Cerqueira C, Soja AMB, Gislason GH, Larsen ML, Andersen UO, Gustafsson I, Thomsen KK, Boye Hansen L, Hammer S, Viggers L, Christensen B, Kvist B, Lindström Egholm C, May O. The Danish cardiac rehabilitation database. *Clin Epidemiol* 2016;**8**:451–456.
30. Braun V, Clarke, V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;**3**:77–101.
31. Christensen AV, Dixon JK, Juel K, Ekholm O, Rasmussen TB, Borregaard B, Mols RE, Thyrsøe L, Thorup CB, Berg SK. Psychometric properties of the Danish Hospital Anxiety and Depression Scale in patients with cardiac disease: results from the DenHeart survey. *Health Qual Life Outcomes* 2020;**18**:9.
32. Rickham PP. Human experimentation. Code of ethics of the World Medical Association. Declaration of Helsinki. *Br Med J* 1964;**2**:177.
33. Lemay KR, Tulloch HE, Pipe AL, Reed JL. Establishing the minimal clinically important difference for the hospital anxiety and depression scale in patients with cardiovascular disease. *J Cardiopulm Rehabil Prev* 2019;**39**:E6–E11.
34. Enrique A, Palacios JE, Ryan H, Richards D. Exploring the relationship between usage and outcomes of an internet-based intervention for individuals with depressive symptoms: Secondary analysis of data from a randomized controlled trial. *J Med Internet Res* 2019;**21**:e12775.
35. Lundgren J, Andersson G, Dahlström Ö, Jaarsma T, Köhler AK, Johansson P. Internet-based cognitive behavior therapy for patients with heart failure and depressive symptoms: a proof of concept study. *Patient Educ Couns* 2015;**98**: 935–942.
36. Wallin E, Norlund F, Olsson EMG, Burell G, Held C, Carlsson T. Treatment activity, user satisfaction, and experienced usability of internet-based cognitive behavioral therapy for adults with depression and anxiety after a myocardial infarction: Mixed-methods study. *J Med Internet Res* 2018;**20**:e87.
37. Lundgren J, Johansson P, Jaarsma T, Andersson G, Kärrer Köhler A. Patient experiences of web-based cognitive behavioral therapy for heart failure and depression: qualitative study. *J Med Internet Res* 2018;**20**:e10302.
38. Titov N, Dear B, Niessen O, Staples L, Hadjistavropoulos H, Nugent M, Adlam K, Nordgreen T, Bruvik KH, Hovland A, Repål A, Mathiasen K, Kraepelien M, Blom K, Svanborg C, Lindefors N, Kalso V. ICBT in routine care: a descriptive analysis of successful clinics in five countries. *Internet Interv* 2018;**13**:108–115.
39. Egholm CL, Helmark C, Doherty P, Nilsen P, Zwisler AD, Bunkenborg G. “Struggling with practices”—a qualitative study of factors influencing the implementation of clinical quality registries for cardiac rehabilitation in England and Denmark. *BMC Health Serv Res* 2019;**19**: 102.
40. Neubeck L, Hansen T, Jaarsma T, Klompstra L, Gallagher R. Delivering healthcare remotely to cardiovascular patients during COVID-19: a rapid review of the evidence. *Eur J Cardiovasc Nurs* 2020;**19**:486–494.