

Angiogenesis inhibitors for metastatic colorectal cancer

Pfeiffer, Per; Liposits, Gabor; Tarpgaard, Line Schmidt

Published in:
Translational Cancer Research

DOI:
10.21037/tcr-23-1568

Publication date:
2023

Document version:
Final published version

Document license:
CC BY-NC-ND

Citation for pulished version (APA):
Pfeiffer, P., Liposits, G., & Tarpgaard, L. S. (2023). Angiogenesis inhibitors for metastatic colorectal cancer. *Translational Cancer Research, 12*(12), 3241-3244. <https://doi.org/10.21037/tcr-23-1568>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.
Unless otherwise specified it has been shared according to the terms for self-archiving.
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.
Please direct all enquiries to puresupport@bib.sdu.dk



Angiogenesis inhibitors for metastatic colorectal cancer

Per Pfeiffer^{1,2^}, Gabor Liposits^{2,3^}, Line Schmidt Taarpgaard^{1,2^}

¹Department of Oncology, Odense University Hospital, Odense, Denmark; ²Department of Clinical Research, University of Southern Denmark, Odense, Denmark; ³Department of Oncology, Gødstrup Hospital, Herning, Denmark

Correspondence to: Per Pfeiffer, MD, PhD. Department of Oncology, Odense University Hospital, Sdr. Boulevard 29, Odense 5000, Odense, Denmark. Department of Clinical Research, University of Southern Denmark, Odense, Denmark. Email: per.pfeiffer@rsyd.dk.

Comment on: Dasari A, Lonardi S, Garcia-Carbonero R, *et al.* Fruquintinib versus placebo in patients with refractory metastatic colorectal cancer (FRESCO-2): an international, multicentre, randomised, double-blind, phase 3 study. *Lancet* 2023;402:41-53.

Keywords: Colorectal cancer (CRC); angiogenesis inhibitors; fruquintinib

Submitted Aug 29, 2023. Accepted for publication Oct 30, 2023. Published online Dec 26, 2023.

doi: 10.21037/tcr-23-1568

View this article at: <https://dx.doi.org/10.21037/tcr-23-1568>

With the recent publications of two important studies, FRESCO-2 (1) and SUNLIGHT (2), we now have data from phase 3 randomized trials showing benefit of anti-angiogenic therapy across the continuum of care, from first to fourth line, in patients with metastatic colorectal cancer (mCRC).

Colorectal cancer (CRC) is the third most prevalent cancer worldwide. Every year around 1.9 million new patients were diagnosed. Approximately half of the patients will, sooner or later, be diagnosed with mCRC, either at the time of diagnosis or later, due to recurrence. Unfortunately, more than 0.9 million (almost 50% of CRC patients) die each year posing a significant challenge for the oncologist community (3).

Since the turn of the millennium, therapy of patients with mCRC has changed extensively, and a doublet or triplet combination of chemotherapy with a targeted agent is now widely used (4,5). Modern treatment strategy for patients with mCRC consists of sequential lines of different systemic therapies. However, with each treatment line, 20–50% of patients will not be candidates for further therapy. The proportion depends very much on patient selection; patients included in clinical trials more often receive further lines of therapy than those in unselected cohorts. The European Society for Medical Oncology has stated that the treatment goal is that 50% of ‘fit’ first-line patients should receive

third-line therapy (4), however, in real-world cohorts only 25–30% of ‘real-world first-line patients’ are receiving third-line therapy and only 10–15% fourth-line (6,7).

First-line chemotherapy consists of a combination of 5-fluorouracil (5FU) and irinotecan followed upon progression by 5FU and oxaliplatin (or vice versa) and often a molecular-targeted drugs will be added, e.g., epidermal growth factor receptor inhibitors (for selected patients whose tumors are *RAS*^{wt}/*BRAF*^{wt}) or anti-angiogenic drugs which can be used in all patients independently of the molecular subtypes (4,5).

The treatment of patients with mCRC, has in general changed from ‘one strategy fits all’ to a more personalized approach. This strategy is, however, not yet the case for anti-angiogenic therapy due to the absence of validated predictive biomarkers. In this commentary, we will stick to the therapeutic approach in molecularly unselected patients and leave the discussion of ‘druggable alterations’ to others (5,8).

Angiogenesis refers to the creation of new blood vessels from already present endothelial cells. Angiogenesis plays an important part in the continued growth of malignant cells since growth beyond a few millimeters requires a blood supply (9). However, malignant blood vessels are often functionally abnormal and consist of an unbalanced network of leaky blood vessels, which result in a high intratumoral

[^] ORCID: Per Pfeiffer, 0000-0002-2925-0586; Gabor Liposits, 0000-0002-8204-3949; Line Schmidt Tarpgaard, 0000-0001-8505-2544.

pressure. Folkman (10) described this dependency 50 years ago and the factor that stimulated the formation of new blood vessels was later named vascular endothelial growth factor (VEGF). This system comprises of 6 ligands and 3 receptors (VEGFR).

Targeting angiogenesis represents a key element in the overall treatment strategy and angiogenesis inhibition is in general clinically effective and has an important place in the therapeutic armamentarium of mCRC. Anti-angiogenic drugs consist of 2 main groups (9); the monoclonal antibodies (mAbs) and the small molecules, tyrosine kinase inhibitors (TKIs). The mAbs bind directly to VEGF-A or block the extracellular binding domain of the matching receptor.

Bevacizumab, a humanized mAb that targets VEGF-A and was approved in 2004, is used in combination with chemotherapy in first-line or in second-line treatment or, according to a 'beyond progression' strategy, in both first- and second-line with a change of the chemotherapy backbone. Two other anti-angiogenic drugs (aflibercept, a recombinant fusion protein and ramucirumab, a fully human mAb that exclusively targets the VEGFR-2 ligand-binding domain) are approved in combination with chemotherapy as second-line treatment (9). As monotherapy, mAbs have limited efficacy in mCRC, but in general, anti-angiogenic therapy increases efficacy of chemotherapy with a slightly additional toxicity (9).

The small molecules, TKIs, exert their anti-angiogenic effect after internalization in the cell and binding to, and inhibit the kinase domain of the various receptors involved in the angiogenesis. Despite many studies including thousands of patients—in contrast to the mAbs—no randomized study has shown a survival benefit of TKIs in combination with chemotherapy neither in first- or second-line therapy in mCRC (9). So far, only regorafenib has been implemented in the treatment sequence of mCRC patients.

As first-line therapy, bevacizumab in combination with 5FU monotherapy or IFL (an outmoded bolus regimen combining irinotecan and 5FU), substantially improved efficacy with a median overall survival (OS) benefit of 3–6 months (9). However, in combination with modern regimens (e.g., FOLFOX or CapOx), the advantage was more modest and the survival benefit was only around 1.5 months (9,11,12). As second-line therapy with non-cross resistant chemotherapy, bevacizumab (or aflibercept/ or ramucirumab) prolonged median survival with about two months, also in patients who had received bevacizumab as part of first-line therapy.

Until recently, after failure of the first two subsequent lines of therapy, treatment possibilities for patients with molecularly unselected mCRC were limited to two orally administered drugs: regorafenib (13) and trifluridine–tipiracil (FTD/TPI) (14).

Regorafenib is an oral multikinase inhibitor that blocks numerous protein kinases including VEGFR1. Compared to placebo, regorafenib prolonged median OS with 1.4 months (from 5.0 to 6.4 months). The most common grade 3 or more symptomatic adverse events were hand-foot skin reaction (17%), fatigue (10%), diarrhea (7%), and rash (6%). In 2012, regorafenib was approved as third-line monotherapy in unselected patients with chemo-refractory mCRC.

FTD/TPI consists of an antineoplastic thymidine-based nucleoside analogue, trifluridine, and a thymidine phosphorylase inhibitor, tipiracil hydrochloride, which prevents the rapid breakdown of trifluridine. Compared to placebo, FTD/TPI prolonged median OS with 1.8 months (from 5.3 to 7.1 months) with limited toxicity. In 2015, FTD/TPI was approved as a third-line monotherapy option in unselected patients with chemo-refractory mCRC.

However, angiogenesis received renewed interest after the publication of SUNLIGHT and FRESCO-2 trials, showing benefit of anti-angiogenic therapy across all lines of therapy (1,2).

A Danish randomized trial including 93 patients with mCRC showed a median survival benefit of 2.7 months [from 6.7 to 9.4 months; hazard ratio (HR) =0.55, P=0.028] for the combination of bevacizumab with FTD/TPI compared to FTD/TPI with a comparable benefit across patient subgroups, including prior treatment with bevacizumab in the immediate past line of therapy (15). These promising results were recently confirmed by data from the large SUNLIGHT study which had a similar design.

SUNLIGHT was an international, randomized phase 3 study, including 492 patients with metastatic colorectal adenocarcinoma. Patients should have received a maximum of two prior lines of systemic treatment including all current standard approved therapies (5FU, oxaliplatin, irinotecan, anti-EGFR (only for *RAS*wt), and anti-VEGF mAb). The combination of bevacizumab with FTD/TPI prolonged median OS with 3.3 months from 7.5 to 10.8 months (HR =0.61; P<0.001). The progression-free survival (PFS) was prolonged from 2.4 to 5.6 months in the combination group (HR =0.44; P<0.001). In both groups, most common adverse events were hematological toxicities (neutropenia and anemia), hypertension, and nausea.

Consequently, in June 2023, the European Medicines Agency (EMA) and in August 2023 the U.S. Food and Drug Administration (FDA) adopted a positive opinion recommending a change to the terms of the marketing authorisation for FTD/TPI (Lonsurf) to: Lonsurf is indicated in combination with bevacizumab for the treatment of adult patients with mCRC who have received two prior anticancer treatment regimens including fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapies, anti-VEGF agents, and/or anti-EGFR agents.

The importance of VEGFR is paramount in tumor angiogenesis and progression. The new pan-VEGFR inhibitor fruquintinib was developed to enhance kinase selectivity resulting in improved efficacy and better tolerability. Fruquintinib demonstrated a favorable toxicity profile as monotherapy and is currently investigated in combination with other drugs in clinical trials. Based on data from the FRESCO trial, fruquintinib has been approved in China (16). The FRESCO trial was a Chinese phase 3 trial including 416 patients with chemo-refractory mCRC and patients were randomized to fruquintinib or placebo. Treatment with fruquintinib prolonged median OS with 2.7 months (from 6.6 to 9.3 months, HR =0.65; P<0.001).

FRESCO-2 was an international, randomized, placebo-controlled study conducted in 14 countries and in less than 16 months, including 691 patients with mCRC (1). Patients had received all current standard approved therapies (5FU, oxaliplatin, irinotecan, anti-EGFR (only for *RAS*wt), and bevacizumab) and patients should be resistant or intolerant to FTD/TPI, or regorafenib (or both). FRESCO-2 included more heavily pretreated patients than SUNLIGHT.

The primary endpoint of the FRESCO-2 study, OS was 7.4 months in the fruquintinib arm *vs.* 4.8 months with placebo arm (HR =0.66, 95% CI: 0.55–0.80; P<0.0001). Fruquintinib also improved PFS from a median of 1.8 to 3.7 months (HR =0.32; P<0.0001). At 6 months, 24% of patients in the fruquintinib group versus 1% in the placebo group had no sign of progressive disease. Fruquintinib improved OS and PFS across all patient subgroups, including prior treatment with FTD/TPI or regorafenib. Only 25 patients had not received prior VEGF inhibitors and in this small group the advantage was extensive (HR =0.19), but also prior anti-VEGF treated patients benefitted (HR =0.68). Common symptomatic grade ≥ 3 adverse events were), asthenia (8%) and hand-foot syndrome (6%). Treatment discontinuation due to adverse events occurred

in 20% and 21% of the patients receiving fruquintinib *vs.* placebo, respectively.

Currently, a New Drug Application for fruquintinib undergoes priority review by the FDA for mCRC indication in line with the FRESCO-2 inclusion criteria.

As third line therapy in unselected patients with mCRC, the combination of FTD/TPI with bevacizumab will be the new standard of care in mCRC and fruquintinib will probably be the standard of care in patients progressing to FTD/TPI and bevacizumab even though only few patients in FRESCO-2 study did receive the FTD/TPI and bevacizumab combination immediately prior starting fruquintinib. In the FRESCO-2 study, all subgroups seemed to benefit from fruquintinib but it would be interesting to see if patients receiving bevacizumab in the immediate prior line before starting fruquintinib do benefit to the same extent from this new fascinating anti-angiogenic therapy.

The results of the SUNLIGHT and the FRESCO-2 studies support the use of anti-angiogenic therapy in the continuum of care for patients with chemo-refractory mCRC. A significant number of ongoing and planned phase 2 and 3 trials, mainly Chinese, are testing promising and interesting combinations of fruquintinib with chemotherapy (including FTD/TPI), targeted agents, and immunotherapy, also in earlier lines of therapy. Results from these trials will hopefully guide to the best clinical practice and establish the place of emerging treatment options in the continuum of care for patients with mCRC.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Translational Cancer Research*. The article has undergone external peer review.

Peer Review File: Available at <https://tcr.amegroups.com/article/view/10.21037/tcr-23-1568/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://tcr.amegroups.com/article/view/10.21037/tcr-23-1568/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all

aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Dasari A, Lonardi S, Garcia-Carbonero R, et al. Fruquintinib versus placebo in patients with refractory metastatic colorectal cancer (FRESCO-2): an international, multicentre, randomised, double-blind, phase 3 study. *Lancet* 2023;402:41-53.
2. Prager GW, Taieb J, Fakih M, et al. Trifluridine-Tipiracil and Bevacizumab in Refractory Metastatic Colorectal Cancer. *N Engl J Med* 2023;388:1657-67.
3. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021;71:209-49.
4. Van Cutsem E, Cervantes A, Adam R, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016;27:1386-422.
5. Cervantes A, Adam R, Roselló S, et al. Metastatic colorectal cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol* 2023;34:10-32.
6. Sorbye H, Dragomir A, Sundström M, et al. High BRAF Mutation Frequency and Marked Survival Differences in Subgroups According to KRAS/BRAF Mutation Status and Tumor Tissue Availability in a Prospective Population-Based Metastatic Colorectal Cancer Cohort. *PLoS One* 2015;10:e0131046.
7. Abrams TA, Meyer G, Schrag D, et al. Chemotherapy usage patterns in a US-wide cohort of patients with metastatic colorectal cancer. *J Natl Cancer Inst* 2014;106:djt371.
8. Avagyan S, Binder H. Subtyping or not subtyping-Quo vadis for precision medicine of colorectal cancer. *Transl Cancer Res* 2023;12:1067-72.
9. Folkman J. Tumor angiogenesis: therapeutic implications. *N Engl J Med* 1971;285:1182-6.
10. Hansen TF, Qvortrup C, Pfeiffer P. Angiogenesis Inhibitors for Colorectal Cancer. A Review of the Clinical Data. *Cancers (Basel)* 2021;13:1031.
11. Botrel TEA, Clark LGO, Paladini L, et al. Efficacy and safety of bevacizumab plus chemotherapy compared to chemotherapy alone in previously untreated advanced or metastatic colorectal cancer: a systematic review and meta-analysis. *BMC Cancer* 2016;16:677.
12. Baraniskin A, Buchberger B, Pox C, et al. Efficacy of bevacizumab in first-line treatment of metastatic colorectal cancer: A systematic review and meta-analysis. *Eur J Cancer* 2019;106:37-44.
13. Grothey A, Van Cutsem E, Sobrero A, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013;381:303-12.
14. Mayer RJ, Van Cutsem E, Falcone A, et al. Randomized trial of TAS-102 for refractory metastatic colorectal cancer. *N Engl J Med* 2015;372:1909-19.
15. Pfeiffer P, Yilmaz M, Möller S, et al. TAS-102 with or without bevacizumab in patients with chemorefractory metastatic colorectal cancer: an investigator-initiated, open-label, randomised, phase 2 trial. *Lancet Oncol* 2020;21:412-20.
16. Li J, Qin S, Xu RH, et al. Effect of Fruquintinib vs Placebo on Overall Survival in Patients With Previously Treated Metastatic Colorectal Cancer: The FRESCO Randomized Clinical Trial. *JAMA* 2018;319:2486-96.

Cite this article as: Pfeiffer P, Liposits G, Taarpgaard LS. Angiogenesis inhibitors for metastatic colorectal cancer. *Transl Cancer Res* 2023;12(12):3241-3244. doi: 10.21037/tcr-23-1568